

Modern Concepts of Cardiovascular Disease

Published monthly by the AMERICAN HEART ASSOCIATION

50 WEST 50TH STREET, NEW YORK, N. Y.

DR. WILLIAM J. KERR, San Francisco, Editor

DR. JOHN J. SAMPSON, San Francisco, Associate Editor

85228
1

Vol. VI

February, 1937

No. 2

THE CARDIAC CRIPPLE IN INDUSTRY

This is a problem which is complicated by two independent and generally conflicting desiderata—the interest of the patient and the interest of employers and others.

The interest of the patient calls for the best possible understanding of his condition, from this the optimal conception of his capacity and safety in various kinds of work, and then his placement in the most suitable employment. Rules of thumb, cross referring from pathological heart conditions to kinds of work permissible, would be convenient, but would also be unreliable. The same would be true of a rigid classification based on functional tests. The need is for a broad and balanced view of all possible elements in the individual situation: the heart lesion (and if originally infectious, the liability to reinfection), the age, habitus, general bodily condition, nervous-mental constitution, training, and even the predilections of the patient; and the kind and hours of work, ease of travel to and from work, climate and altitude, and opportunities for proper nourishment, sleep, recreation—and equanimity.

Active infection of the heart usually precludes any recommendation of employment; and it should be assumed to exist at least for several months following the fever or other obvious clinical manifestations of rheumatic fever, chorea, or other infection known or presumed to have involved the heart. And beyond this period of convalescence the prudent physician, knowing the protracted and insidious habit of the infection, will commend to the young person with rheumatic heart disease only such work as contributes to or at least is consistent with maintenance of the best possible condition of the body to resist it. In lieu the general bodily condition, though important to maintain, is not the chief therapeutic objective; and active cardiovascular syphilis is compatible with suitable work during the period of treatment if liberal cardiac reserve strength is present.

When active infection is not an issue, the fitness of the heart should be judged by an appraisal of the present virtues and probable future of the myocardium—the stability of its mechanism of beat, the adequacy of its blood supply, and its margin of ability to carry peak loads safely with existing peripheral, valvular or other handicap.

Overload—Hypertrophy—Dilatation—Sequence. This is the gradual progress of the handicapped myocardium toward congestive failure. Contrary to a now prevalent notion, accurate knowledge of valve lesions is important, indeed is prerequisite to a full

understanding of the myocardium. Illustration: A young man with healed rheumatic endocarditis, moderate cardiac enlargement, regular heart rhythm, and ability to climb stairs comfortably. If he has exclusively or predominantly mitral disease—let us say, insufficiency and beginning stenosis—the outlook is for slowly advancing stenosis, eventual paroxysmal or continuous atrial fibrillation, and, without adequate precautions, ultimately a zigzag downward course of lapses into and partial recoveries from congestive failure. Meantime his cardiac reserve can be safeguarded by a relatively simple prescription of employment—to avoid exercise which causes any dyspnea; for in his case the right ventricle carries the extra load (at the other end of the pulmonary circuit), and its embarrassment is immediately reflected in engorgement of lung vessels and commensurate diminution of air space. If, however, his valve lesion is exclusively or predominantly aortic, the outlook without intelligent management is for gradual increase in heart size with little or no discomfort and considerable tolerance for exercise until the ultimate, and then often irremediable, failure. To give him the same simple injunction, “avoid exercise which causes hard breathing,” betrays a deplorable misconception of his myocardial situation; for his pulmonary circuit, protected by a competent mitral valve, is likely not to be much engorged until his left ventricle, greatly hypertrophied, has expended its reserve and dilates beyond its ability to recuperate. His exercise never should be strenuous, never more than mildly fatiguing. Prior to the onset of incipient failure (when almost any employment would be hazardous) the best single guide for his work prescription is the size of the heart; but this should be viewed in its relation to the duration of the valvulitis, the severity of the lesion (character of aortic second sound and peripheral phenomena), the age, build and previous activities of the patient.

In varying degree similar concepts of myocardial situation apply to other overloads which tend to advance or retard the appearance of dyspnea. Like mitral stenosis, pulmonary emphysema by a somewhat different mechanism gives an early signal of embarrassment of the right ventricle. Like aortic disease, hyperpiesis, coarctation of the aorta, thyrotoxicosis and arteriovenous aneurysm overwork the left ventricle and do not induce much dyspnea until it is far advanced in the hypertrophy-dilatation sequence. Pericardial adhesions and combinations of valvular or other handicaps usually present less

ANNOUNCEMENT!

For the benefit of those who wish to save current issues of the “Modern Concepts of Cardiovascular Disease,” we have had designed a handsome loose leaf binder, in green fabrikoid with gold stamping, which will accommodate fifty issues. The price is \$1.00, postpaid.

There is still available some copies of the bound volume containing the issues of 1932 to 1935 inclusive, with subject and author indices, price \$2.00, postpaid.

Orders with remittances should be mailed to the American Heart Association, 50 West 50th Street, New York, N. Y.

clearly differentiated relationships between dyspnea and the hypertrophy-dilatation sequence; but in estimating a patient's position in this sequence, i.e., his cardiac reserve and his optimal exercise, the heart size is always of great importance.

Myocardial Nutrition Deficit. To illustrate further the importance of studying valve condition: If our hypothetical patient is middle aged when he begins to experience cardiac symptoms, and we then find aortic insufficiency exclusively, we should realize that in all probability he has syphilis (regardless of history and Wassermann test), and that lesions around the valve may be encroaching on the mouths of the coronary arteries. Here a view of the myocardial situation limited to the hypertrophy-dilatation sequence is likely to be disastrously inadequate. More important is a conception of the threat to heart muscle nutrition, the doubtful possibility that antisyphilitic treatment will preserve it, and meantime the danger of an irreversible downward course, of syncopal attacks, of angina or sudden death from block, ventricular fibrillation or other disorder of heart-beat mechanism.

Coronary sclerosis advances insidiously, often to the danger point before it can be detected even by electrocardiography. Mild substernal distress with effort or a little nocturnal dyspnea are common early indications. Fully developed angina pectoris need not be discussed here, except to urge meticulous history-taking as superior to all other forms of examination—not only for diagnosis but also and especially for guidance in prescribing work. Muscular effort or mental excitement, independently or in relation to meals, temperature, hour of the day, etc., and anything that causes attacks must be avoided. Individualized advice is essential. The same is true after recovery from acute coronary occlusion. Beginning after sufficient time for firm scarring of the infarct, exercise may be gradually and cautiously increased. If the heart is not enlarged and gives no indication of embarrassment, heavy work sometimes can be resumed with impunity, although as a rule slow or stop signals appear long before this point is reached. Recurrence is common with or without work.

All persons with myocardial nutrition deficit, either absolute (as in acute coronary occlusion), relative (as in effort-induced angina), or potential (as in aortic syphilis which threatens the coronary mouths) are liable to syncope or sudden death through ventricular fibrillation or other abnormal beat mechanism. The danger diminishes as time elapses after a coronary accident or after the beginning of successful antisyphilitic treatment; but it is never quite absent, and requires careful consideration in the case of would-be chauffeurs, railway engineers and many others. And, (of course, *cerebral nutrition deficit* also should be remembered—the vascular accidents, transient vertigo, or simple blank moments common with high blood pressure, postural syncope with low pressure, reflex syncope, etc.)

Myocarditis. True infection of heart muscle, either acute or chronic, and *myocardial fibrosis* or other forms of degeneration are usually advanced when clinically recognizable and require much conservatism in prescribing work. With low reserve power and poor material for building reserves, compensation is easily broken and hard to restore. The graver forms of abnormal beat mechanism are common, and may induce syncopal attacks with special hazards to the patient, to others and to property. The history of such attacks, objective evidence of muscle damage, (especially of the conduction system), pain, dyspnea or exhaustion with little provocation, heart enlargement, and congestive phenomena are important data, in varying degrees and combinations contributing to sound judgment about employment possibilities.

Masquerade Heart Disease. Inconsequential systolic murmurs still bar the way of many applicants for employment (and such rejections too often inflict further and more serious injury by implanting anxiety neurosis). To some extent the same is true

of simple ventricular ectopic beats and other benign arrhythmias. Paroxysmal tachycardia and even atrial fibrillation may exist without demonstrable structural heart disease and without important reduction in cardiac efficiency. In general, while systolic murmurs and arrhythmias certainly cannot be ignored, they should take second place in the mind of the examiner; and for practical purposes he should pass as normal many hearts with these peculiarities, if other and more important data warrant it, especially the size of the heart and comfortable tolerance of exercise.

But tolerance of exercise requires discriminating interpretation. It may be much reduced without real heart disease, as in anemia, respiratory diseases, certain glandular disturbances, fatigue states, etc. Judgment then should be based upon the underlying condition rather than upon the heart, which acts merely as spokesman. Patients with neurocirculatory asthenia may be quite unfit for much exertion; but for them to realize that the necessary exercise restriction is designed for nervous comfort rather than cardiac safety is decidedly to their advantage and that of their work.

The interest of employers calls for efficient and safe service. But efficiency is not merely the ability to do certain work. Speed, maintained through regular working hours, and freedom from much sick-leave are usually required. And safety—that is not simply immunity from syncopal attacks for machine and transportation operatives and the like. Important (and often astonishingly neglected) as this is, it is less in the minds of many employers than their own immunity from penalties under the industrial accident laws. The administration of these laws often gives employees advantages far beyond equitable compensation for bona fide industrial injuries. A competent machinist with mitral stenosis may be crippled rather suddenly by the onset of atrial fibrillation. The chances are that some very ordinary act in his work will be blamed for it, that his attorney will find medical testimony to support a claim, and that the man will retire for an indefinite time at the expense of the employer. Or an aneurysm bursts in the act of lifting a typewriter; the lift has no important effect on the duration of the man's life—merely a "last straw" in the causative chain of events—and yet the widow receives an award, the same award as if her husband had been well and had been blown to bits by a boiler explosion. All very good for this particular machinist and this widow, but hard on the next applicant for employment with mitral stenosis or aneurysm. Physically and otherwise he may be well qualified for the job he seeks. He may be anxious to assume all risk incident to his disease. Will his signed waiver protect the employer? Very doubtfully so, advise the lawyers. Can the employer protect himself by insurance? Yes, but eventually more "accidents" will bring higher premiums.

And finally, opportunity for the cardiac cripple is often restricted by fellow workers. Their hospital association may be burdened by expense if he is employed. Even the old employee may feel the pinch of the group. A locomotive engineer recovers well after coronary thrombosis, but cannot safely run his engine again. He is qualified for clerical work and the railway management would like to give it to him, but is restrained by the clerks' union, who will brook no trespassing on their seniority.

The problem of employment for cardiac cripples is expanding in both its parts. With longer average life, degenerative diseases increase the number of cripples; and with present social and economic trends their opportunities are more and more curtailed. The medical part of the problem has received much attention, the social-economic part very little. The latter involves a complicated web of group forces opposed to the interest of the cardiac patient, and will require collective action for his relief.

Eugene S. Kilgore, M.D.

San Francisco, California

1
 2
 3
 4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525